

Characterisation of *rpsL* and *rrs* mutations associated with streptomycin resistance in multidrug-resistant *Mycobacterium tuberculosis* clinical isolates (from Poland)

Zofia Bakula¹, Helena Ignatowska¹, Katarzyna Roeske¹, Michał Kamiński¹, Agnieszka Napiórkowska², Jacek Bielecki¹, Ewa Augustynowicz-Kopec², and Tomasz Jagielski¹

¹Department of Applied Microbiology, Institute of Microbiology, Faculty of Biology, University of Warsaw, Miecznikowa 1, 02-096 Warsaw, Poland

²Department of Microbiology, National Tuberculosis and Lung Diseases Research Institute, Warsaw, Płocka 26, 01-138 Warsaw, Poland

The study was financed by the Ministry of Science and Higher Education: «Juventus Plus» grant (Project No. IP2011018771) and the Faculty of Biology, University of Warsaw intramural grant DSM.

Background

Streptomycin (SM) is a component of the first-line regimens for the treatment of tuberculosis that acts by binding to the 30S subunit of the ribosome, and thus inhibiting protein synthesis [1]. Mutations in genes *rpsL* and *rrs* encoding the ribosomal protein S12, and the 16S rRNA, respectively, have been associated with resistance to SM in *Mycobacterium tuberculosis* clinical isolates [2, 3]. Analysis of mutations in the *rpsL* and *rrs* genes in *M. tuberculosis* may contribute to the development of new diagnostic tests allowing for rapid detection of SM resistance.

Objective

The aim of this study was to investigate the presence of mutations in two genetic loci, namely the *rpsL* and *rrs* genes, most frequently associated with SM resistance.

Methods

A total of 50 *M. tuberculosis* (32 SM-resistant and 18 SM-susceptible) strains isolated from as many MDR-TB patients in Poland throughout 2004 were included in the study. Mutations in the *rpsL* and *rrs* genes were detected through PCR amplification of both loci, followed by their direct sequencing in forward and reverse directions. The obtained nucleotide sequences were aligned against the reference sequences of the respective genes of *M. tuberculosis* reference strain H₃₇Rv. The obtained results were interpreted in the context of SM susceptibility profile of the strains tested. Primary isolation, species identification, and drug susceptibility testing had been done beforehand, as described elsewhere [4].

Results

Among the 32 SM-resistant isolates, 24 (75%) contained mutations in either the *rpsL* (16 isolates) or the *rrs* (8) gene. None of the isolates had mutations in both loci concurrently. Of the 16 SM-resistant isolates with *rpsL* gene mutations, 14 (87.5%) isolates had a substitution in codon 43 from lysine to arginine, whereas one isolate had the same amino acid substitution in codon 88. These two mutation types were found exclusively in SM-resistant strains. One SM-resistant isolate had a silent mutation in codon 39. This mutation was also found in two SM-susceptible isolates. Four *rrs* mutation types were identified. They occurred at positions 517 in 4 isolates, 906 in two isolates, 514 in one isolate, and 907 in another isolate. All isolates with mutations in the *rrs* gene were SM-resistant. Eight (25%) SM-resistant isolates were wild-type for both loci analysed.

Conclusions

The study revealed a relatively high frequency of mutations in SM resistance-related genes. Every nonsynonymous mutation detected in the *rpsL* or *rrs* gene was found exclusively in SM-resistant isolates. This underlines the usefulness of *rpsL* and *rrs* mutations as molecular markers predictive of resistance of tubercle bacilli to SM.

Keywords: *Mycobacterium tuberculosis*; *rpsL*; *rrs*; streptomycin; multidrug-resistance; sequence analysis

References

- [1] Cole, S. T., and A. Telenti. 1995. Drug resistance in *Mycobacterium tuberculosis*. Eur. Respir. J. Suppl. **20**:701-713.
- [2] Finken, M., P. Krischner, A. Meier, A. Wrede, and E. C. Böttger. 1993. Molecular basis of streptomycin resistance in *Mycobacterium tuberculosis*: alterations of the ribosomal protein S12 gene and point mutations within a functional 16S ribosomal RNA pseudoknot. Mol. Microbiol. **9**:1239-1246.

- [3] Tracevska, T., I. Jansone, A. Nodieva, O. Marga, G. Skenders, V. Baumanis. 2004. Characterisation of *rpsL*, *rrs* and *embB* mutations associated with streptomycin and ethambutol resistance in *Mycobacterium tuberculosis*. Res. Microbiol., 155:830-834.
- [4] Augustynowicz-Kopeć, E., Z. Zwolska, A. Jaworski, E. Kostrzewa, M. Klatt. 2003. Drug-resistant tuberculosis in Poland in 2000: second national survey and comparison with the 1997 survey. Int. J. Tuberc. Lung. Dis., 7:645-651.